

Finite size corrections in the random assignment problem

Sergio Caracciolo,^{*} Matteo P. D'Achille,[†] and Enrico M. Malatesta[‡]
 Dipartimento di Fisica, University of Milan and INFN, via Celoria 16, I-20133 Milan, Italy

Gabriele Sicuro[§]
 Dipartimento di Fisica, Università "Sapienza", P.le A. Moro 2, I-00185, Rome, Italy
 (Dated: February 20, 2017)

We analytically derive, in the context of the replica formalism, the first finite size corrections to the average optimal cost in the random assignment problem for a quite generic distribution law for the costs. We show that, when moving from a power law distribution to gamma distribution, the leading correction changes both in sign and in its scaling properties. We also examine the behavior of the corrections when approaching a delta function distribution. By using a numerical solution of the saddle-point equations, we provide predictions which are very well confirmed by numerical simulations.

I. INTRODUCTION

Matching is a classical problem in combinatorial optimization [1, 2]. It can be defined on any graph $\mathcal{G} = (\mathcal{V}, \mathcal{E})$, where \mathcal{V} is its set of vertices and \mathcal{E} its set of edges. A matching in \mathcal{G} is a set of non-adjacent edges of \mathcal{G} , that is, a set of edges that don't share a common vertex. A matching is *maximal* when the addition of any new edge to the set makes it no longer a matching. A matching is said to be *maximum* when it has the maximal cardinality among the maximal matchings. All the maximum matchings have the same cardinality $\nu(\mathcal{G})$ which is called the *matching number* of the graph \mathcal{G} . A *perfect matching* (or *1-factor*) is a matching which matches all vertices of the graph. That is, every vertex of the graph is incident to exactly one edge of the matching. Every perfect matching is maximum and hence maximal. A perfect matching is also a minimum-size edge cover. We will denote by \mathcal{M} the set of perfect matchings.

Suppose now that we can assign a cost $w_e \geq 0$ to each edge $e \in \mathcal{E}$. For each perfect matching $\pi \in \mathcal{M}$ we define a total cost (energy)

$$E(\pi) := \sum_{e \in \pi} w_e, \quad (1)$$

and a mean cost for edge

$$\epsilon(\pi) := \frac{1}{\nu(\mathcal{G})} \sum_{e \in \pi} w_e. \quad (2)$$

The *matching problem* consists in the choice of the optimal perfect matching $\pi^* \in \mathcal{M}$ which minimizes the total cost

$$E(\pi^*) = \min_{\pi \in \mathcal{M}} E(\pi). \quad (3)$$

Let us now consider the matching problem on the complete bipartite graph $\mathcal{K}_{N,M}$, in which the vertex set is the union of

two disjoint sets \mathcal{V}_1 and \mathcal{V}_2 of cardinality N and M respectively. Denoting by $[n] := \{1, \dots, n\}$, we identify \mathcal{V}_1 and \mathcal{V}_2 with $[N]$ and $[M]$ respectively, and the edge set is therefore the set of all couples $e = (i, j)$ with $i \in [N]$ and $j \in [M]$. In this case the matching problem is usually called the (general) *assignment problem* and the matching number is $\nu(\mathcal{K}_{N,M}) = \min\{N, M\}$. In the following we shall concentrate on the $N = M$ case, in which a perfect matching π is a permutation in the symmetric group \mathcal{S}_N , and can be represented by a square matrix with entries $\pi_{ij} \in \{0, 1\}$ for all $i \in [N]$ and $j \in [N]$, such that

$$\pi_{ij} = \begin{cases} 1 & \text{if } e = (i, j) \in \pi, \\ 0 & \text{otherwise,} \end{cases} \quad (4)$$

with the constraints

$$\sum_{i=1}^N \pi_{ij} = \sum_{j=1}^N \pi_{ji} = 1 \quad \forall j \in [N]. \quad (5)$$

The matching cost associated to π can be written as

$$E(\pi) = \sum_{i=1}^N \sum_{j=1}^N \pi_{ij} w_{ij}. \quad (6)$$

From the point of view of computational complexity, matching problems are *simple* problems, being in the P complexity class, as Kuhn [3] proved in 1955 with his celebrated Hungarian algorithm for the assignment problem. Very fast algorithms are nowadays available both to find perfect matchings and to solve the matching problem on a generic graph [2, 4–6].

The properties of the solution of a matching problem on a given ensemble of realizations is often interesting as well [7]. In the *random assignment problem*, for example, we consider the matching problem on $\mathcal{K}_{N,N}$, whereas the costs for all edges are independent random variables, identically distributed with a common law ρ . Each particular choice $\mathcal{W} = \{w_e\}_{e \in \mathcal{E}}$ for the set of edge costs is called an *instance* of the problem. In this random version of the problem, we are interested in the *typical* properties of the optimal matching. In particular we will concentrate in the asymptotic behaviour for large N of the average optimal cost

$$\overline{E} := \overline{E(\pi^*)} = \min_{\pi \in \mathcal{M}} \sum_{i=1}^N \sum_{j=1}^N \pi_{ij} w_{ij}, \quad (7)$$

^{*} sergio.caracciolo@mi.infn.it

[†] matteopietro.dachille@studenti.unimi.it

[‡] enrico.m.malatesta@gmail.com

[§] gabriele.sicuro@roma1.infn.it

where we have denoted by an overall bar the average over all possible instances (i.e., the average over the *disorder*). In the same way, we can consider the matching problem with random weights on the complete graph \mathcal{K}_{2N} , having $2N$ vertices such that each one of them is connected to all the others. We simply call this variation of the problem *random matching problem*. Both the random matching problem and the random assignment problem have been solved by Mézard and Parisi [8] in 1985 by means of the replica trick. The random assignment problem and the random matching problem have been also generalized to the Euclidean case, in which the weights in \mathcal{W} are functions of distances between points associated to the vertices of the graph, randomly generated on a certain Euclidean domain [9–12]. Due to the underlying Euclidean structure, dimensionality plays an important role in the scaling of the optimal cost of random Euclidean matching problems [10, 13], and correlation functions can be introduced and calculated [14, 15]. Euclidean matching problems proved to be deeply connected with Gaussian stochastic processes [14, 16] and with the theory of optimal transport [17]. In the latter context, Ambrosio *et al.* [18] rigorously derived the asymptotic behavior of the average optimal cost for the 2-dimensional random Euclidean assignment problem, previously obtained in Ref. [13] using a proper scaling ansatz. For a recent review on random Euclidean matching problems, see Ref. [19].

Remarkably enough, after the seminal works of Kirkpatrick *et al.* [20], Orland [21], Mézard and Parisi, the application of statistical physics techniques to random optimization problems proved to be extremely successful in the study of the typical properties of the solutions, but also in the development of algorithms to solve a given instance of the problem [22, 23]. In formulating a combinatorial problem as a model in statistical mechanics, an artificial inverse temperature β is introduced to define a Boltzmann weight $\exp(-\beta E)$ for each configuration. Of course, configurations of minimal energy are the only ones to contribute in the limit of infinite β . For example, in the assignment problem, the corresponding partition function for each instance is

$$\begin{aligned} Z[w] &= \sum_{\pi} \left[\prod_{j=1}^N \delta \left(1 - \sum_{i=1}^N \pi_{ij} \right) \delta \left(1 - \sum_{i=1}^N \pi_{ji} \right) \right] e^{-\beta E(\pi)} \\ &= \left[\prod_{i=1}^N \int_0^{2\pi} \frac{e^{i\lambda_i} d\lambda_i}{2\pi} \int_0^{2\pi} \frac{e^{i\mu_i} d\mu_i}{2\pi} \right] \prod_{i,j} [1 + e^{-i(\lambda_j + \mu_i) - \beta w_{ij}}]. \end{aligned} \quad (8)$$

where the “energy” $E(\pi)$ is given by (6). Thermodynamic informations are obtained from the average total free energy

$$\bar{F} := -\frac{\ln Z}{\beta}, \quad (9)$$

$$\bar{E} = \frac{\partial}{\partial \beta} \beta \bar{F}. \quad (10)$$

In this paper we apply the formalism above to the random assignment problem, where the costs of all the edges are taken to be i.i.d. random variables with probability distribution density

$\rho_r(w)$ such that, in the neighborhood of the origin, ρ_r can be written as

$$\rho_r(w) = w^r \sum_{k=0}^{\infty} \eta_k(r) w^k, \quad r > -1. \quad (11)$$

The constraint $r > -1$ is required to guarantee the integrability of the distribution near the origin. By the general analysis performed in Refs. [8, 21], that we will resume in Section II, the average cost, in the asymptotic regime of an infinite number N of couples of matched points, will depend on the power r which appears in Eq. (11) only, aside from a trivial overall rescaling related to η_0 . More precisely, if \bar{E}_r is the average optimal cost obtained using the law ρ_r , then

$$\hat{E}_r = \lim_{N \rightarrow \infty} \frac{1}{N^{\frac{r}{r+1}}} \bar{E}_r = \frac{r+1}{[\eta_0 \Gamma(r+1)]^{\frac{1}{r+1}}} J_r^{(r+1)} \quad (12a)$$

where

$$J_r^{(\alpha)} := \int_{-\infty}^{+\infty} \hat{G}_r(-u) \mathcal{D}_u^{\alpha} \hat{G}_r(u) du \quad (12b)$$

(we will later specify the meaning of the fractional order derivative \mathcal{D}_u^{α}). The function $\hat{G}_r(y)$ is the solution of the integral equation

$$\hat{G}_r(l) = \int_{-l}^{+\infty} \frac{(l+y)^r}{\Gamma(r+1)} e^{-\hat{G}_r(y)} dy \quad (12c)$$

and it is analytically known for $r = 0$ and, after a proper rescaling of its variable, in the $r \rightarrow \infty$ limit.

Our main results concern with the finite size corrections to the average optimal costs, and they will be presented in Section III, extending the classical achievements in Refs. [24, 25]. In particular, we obtain the expansion

$$\hat{E}_r(N) = \hat{E}_r + \sum_{k=1}^{[r]+1} \Delta \hat{F}_r^{(k)} + \Delta \hat{F}_r^T + \Delta \hat{F}_r^F + o\left(\frac{1}{N}\right), \quad (13a)$$

where $[r]$ is the integer part of r , $[r] \leq r < [r] + 1$ (the sum is absent for $r < 0$), and the corrections have the structure

$$\Delta \hat{F}_r^{(k)} = \frac{\Delta \phi_r^{(k)}}{N^{\frac{k}{r+1}}}, \quad r \geq 0, \quad 1 \leq k \leq [r] + 1, \quad (13b)$$

$$\Delta \hat{F}_r^T = -\frac{1}{N} \frac{\Gamma(2r+2) J_r^{(0)}}{(r+1) \eta_0^{\frac{1}{r+1}} [\Gamma(r+1)]^{\frac{2r+3}{r+1}}} \quad (13c)$$

$$\Delta \hat{F}_r^F = -\frac{1}{N} \frac{1}{2 [\eta_0 \Gamma(r+1)]^{\frac{1}{r+1}} J_r^{(r+3)}}, \quad (13d)$$

$\Delta \phi_r^{(k)}$ being independent on N . In particular, for $r > 0$, we have that, provided that $\eta_1 \neq 0$, the first finite size correction is given by

$$\Delta \hat{F}_r^{(1)} = -\frac{\eta_1}{N^{\frac{1}{r+1}} \eta_0 [\eta_0 \Gamma(r+1)]^{\frac{2}{r+1}}} J_r^{(r)}. \quad (13e)$$

In our discussion, we shall consider in particular two probability distribution densities, namely the gamma distribution

$$\rho_r^\Gamma(w) := \frac{w^r e^{-w} \theta(w)}{\Gamma(r+1)}, \quad (14)$$

defined on \mathbb{R}^+ , and the power law distribution

$$\rho_r^P(w) := (r+1)w^r \theta(w) \theta(1-w), \quad (15)$$

defined on the compact interval $[0, 1]$. Observe that, for the distribution ρ_r^Γ , we have

$$\eta_k^\Gamma(r) = \frac{1}{\Gamma(r+1)} \frac{(-1)^k}{k!}, \quad k \geq 0, \quad (16)$$

whereas in the case of ρ_r^P

$$\eta_k^P(r) = (r+1) \delta_{k,0}, \quad k \geq 0. \quad (17)$$

The case $r = 0$ has already been considered by Mézard and Parisi [24], and subsequently revised and corrected by Parisi and Ratiéville [25]. In the case analyzed in their works, the contributions $\Delta \hat{F}_0^{(1)}$, $\Delta \hat{F}_0^T$ and $\Delta \hat{F}_0^F$ are of the same order. This is not true anymore for a generic distribution with $r \neq 0$. As anticipated, a relevant consequence of our evaluation is that, if $\eta_1 \neq 0$, for $r > 0$ the most important correction comes from $\Delta \hat{F}_r^{(1)}$ and scales as $N^{-\frac{1}{r+1}}$. It follows that, in order to extrapolate to the limit of infinite number of points, the best choice for the law for random links (in the sense of the one that provides closer results to the asymptotic regime) is the pure power law ρ_r^P , where only analytic corrections in inverse power of N are present. Such a remark is even more pertinent in the limit when $r \rightarrow \infty$ at fixed number of points, where the corrections $\Delta \hat{F}_r^{(k)}$ become of the same order of the leading term. Indeed, the two limits $r \rightarrow \infty$ and $N \rightarrow \infty$ commute only if the law ρ_r^P is considered.

The paper is organized as follows. In Section II we review, in full generality, the calculation of the replicated partition function of the random assignment problem. In Section III we evaluate the finite size corrections, discussing the different contributions and proving Eqs. (13). In Section IV we evaluate the relevant $r \rightarrow \infty$ case, pointing out the noncommutativity of this limit with the thermodynamic limit. In Section V we provide the numerical values of the necessary integrals and we compare our prediction with a Monte Carlo simulation for different values of r . In Section VI we give our conclusions.

II. THE REPLICATED ACTION

In the present section, we perform a survey of the classical replica computation for the random assignment problem, following the seminal works of Mézard and Parisi [8, 24] (for a slightly different approach see also Ref. [21]), but we don't adopt their choice to replace β with $\beta/2$. As anticipated in Section I, the computation of the average of $\ln Z$ goes through the replica trick [26]

$$\overline{\ln Z} = \lim_{n \rightarrow 0} \frac{\overline{Z^n} - 1}{n}. \quad (18)$$

In other words, in order to compute $\overline{\ln Z}$ we introduce n non-interacting replicas of the initial system, denoted by the index $a \in [n]$. For each $i \in [N]$, $2n$ replicated fields $\{\lambda_i^a, \mu_i^a\}_{a=1, \dots, n}$ appear to impose the constraints in Eq. (5), obtaining

$$Z^n[w] = \left[\prod_{a=1}^n \prod_{i=1}^N \int_0^{2\pi} \frac{d\lambda_i^a}{2\pi} \int_0^{2\pi} \frac{d\mu_i^a}{2\pi} e^{i(\lambda_i^a + \mu_i^a)} \right] \times \\ \times \prod_{i=1}^N \prod_{j=1}^N \prod_{a=1}^n \left[1 + e^{-i(\lambda_j^a + \mu_i^a) - \beta w_{ij}} \right]. \quad (19)$$

Let $\mathcal{P}([n])$ be the set of subsets of the set $[n]$, and for each subset $\alpha \in \mathcal{P}([n])$ let $|\alpha|$ be its cardinality. Then

$$\prod_{a=1}^n \left[1 + e^{-i(\lambda_j^a + \mu_i^a) - \beta w_{ij}} \right] \\ = \sum_{\alpha \in \mathcal{P}([n])} e^{-\beta |\alpha| w_{ij} - i \sum_{a \in \alpha} (\lambda_j^a + \mu_i^a)} \\ = 1 + \sum_{p=1}^n \binom{n}{p} e^{-\beta p w_{ij} - i \sum_{a \in \alpha} (\lambda_j^a + \mu_i^a)}, \quad (20)$$

where we have extracted the contribution from the empty set in the sum, which is 1, and we have partitioned the contribution from each subset of replicas in terms of their cardinality. This expression is suitable for the average on the costs. From the law ρ_r we want to extract the leading term for large β of the contribution of each subset $\alpha \in \mathcal{P}([n])$ with $|\alpha| = p$. In particular, we define

$$g_\alpha \equiv g_p := \int_0^{+\infty} \rho_r(w) e^{-\beta p w} dw. \quad (21)$$

Due to the fact that short links only participate to the optimal configuration, approximating $\rho_r(w) \sim \eta_0 w^r$, we obtain that the minimal cost for each matched vertex is of the order $N^{\frac{1}{r+1}}$ so that the total energy E and the free energy should scale as $N^{\frac{r}{r+1}}$, that is, the limits

$$\lim_{N \rightarrow \infty} \frac{1}{N^{\frac{r}{r+1}}} \overline{F} = \hat{F} \quad (22)$$

$$\lim_{N \rightarrow \infty} \frac{1}{N^{\frac{r}{r+1}}} \overline{E} = \hat{E} \quad (23)$$

are finite. This regime can be obtained by considering in the thermodynamic limit

$$\beta = \hat{\beta} N^{\frac{1}{r+1}} \quad (24)$$

where $\hat{\beta}$ is kept fixed. As a consequence we set

$$\hat{g}_p := N g_p = N \int_0^{+\infty} \rho_r(w) e^{-p \hat{\beta} N^{\frac{1}{r+1}} w} dw \\ = \sum_{k=0}^{+\infty} \frac{1}{N^{\frac{k}{r+1}}} \frac{\eta_k \Gamma(k+r+1)}{(\hat{\beta} p)^{k+r+1}}. \quad (25)$$

The replicated partition function can be written therefore as

$$\begin{aligned} \overline{Z}^n &= \left[\prod_{a=1}^n \prod_{i=1}^N \int_0^{2\pi} \frac{d\lambda_i^a}{2\pi} \int_0^{2\pi} \frac{d\mu_i^a}{2\pi} e^{i(\lambda_i^a + \mu_i^a)} \right] \prod_{i,j}^N \left(1 + \frac{T_{ij}}{N} \right) \\ &= \left[\prod_{a=1}^n \prod_{i=1}^N \int_0^{2\pi} \frac{d\lambda_i^a}{2\pi} \int_0^{2\pi} \frac{d\mu_i^a}{2\pi} e^{i(\lambda_i^a + \mu_i^a)} \right] \times \\ &\quad \times \exp \left[\frac{1}{N} \sum_{i=1}^N \sum_{j=1}^N \left(T_{ij} - \frac{T_{ij}^2}{2N} \right) + o \left(\frac{1}{N^2} \right) \right], \end{aligned} \quad (26)$$

with

$$T_{ij} := \sum'_{\alpha \in \mathcal{P}([n])} \hat{g}_\alpha e^{-i \sum_{a \in \alpha} (\lambda_j^a + \mu_i^a)} \quad (27)$$

where in the sum \sum' on subsets the emptyset is excluded. If we introduce, for each subset $\alpha \in \mathcal{P}([n])$, the quantities

$$\frac{x_\alpha + i y_\alpha}{\sqrt{2}} := \sum_{i=1}^N e^{-i \sum_{a \in \alpha} \lambda_i^a} \quad (28a)$$

$$\frac{x_\alpha - i y_\alpha}{\sqrt{2}} := \sum_{i=1}^N e^{-i \sum_{a \in \alpha} \mu_i^a} \quad (28b)$$

we can write

$$\sum_{i=1}^N \sum_{j=1}^N T_{ij} = \sum'_{\alpha \in \mathcal{P}([n])} \hat{g}_\alpha \frac{x_\alpha^2 + y_\alpha^2}{2}, \quad (29a)$$

$$\sum_{i=1}^N \sum_{j=1}^N T_{ij}^2 = \sum'_{\substack{\alpha, \beta \in \mathcal{P}([n]) \\ \alpha \cap \beta = \emptyset}} \hat{g}_\alpha \hat{g}_\beta \frac{x_{\alpha \cup \beta}^2 + y_{\alpha \cup \beta}^2}{2}, \quad (29b)$$

The condition $\alpha \cap \beta = \emptyset$ has been imposed because there are no contributions proportional to factors $\exp(-2i\lambda_i^a)$ or $\exp(-2i\mu_i^a)$ in the partition function. We perform now a Hubbard–Stratonovich transformation, neglecting $o(N^{-2})$ terms in the exponent in Eq. (26), obtaining

$$\begin{aligned} &\exp \left[\frac{1}{N} \sum_{i=1}^N \sum_{j=1}^N \left(T_{ij} - \frac{T_{ij}^2}{2N} \right) \right] = \\ &= \left[\prod'_{\alpha \in \mathcal{P}([n])} \iint \frac{N dX_\alpha dY_\alpha}{2\pi \hat{g}_\alpha} \exp(x_\alpha X_\alpha + y_\alpha Y_\alpha) \right] \times \\ &\times \exp \left[-N \sum'_{\alpha \in \mathcal{P}([n])} \frac{X_\alpha^2 + Y_\alpha^2}{2\hat{g}_\alpha} - \sum'_{\substack{\alpha, \beta \in \mathcal{P}([n]) \\ \alpha \cap \beta = \emptyset}} \hat{g}_\alpha \hat{g}_\beta \frac{X_{\alpha \cup \beta}^2 + Y_{\alpha \cup \beta}^2}{g_{\alpha \cup \beta}^2 4N} \right], \end{aligned} \quad (30)$$

up to higher order terms in the exponent. Now, let us observe that

$$x_\alpha X_\alpha + y_\alpha Y_\alpha = \left(\sum_{i=1}^N e^{-i \sum_{a \in \alpha} \lambda_i^a} \right) \frac{X_\alpha - i Y_\alpha}{\sqrt{2}}$$

$$+ \left(\sum_{i=1}^N e^{-i \sum_{a \in \alpha} \mu_i^a} \right) \frac{X_\alpha + i Y_\alpha}{\sqrt{2}} \quad (31)$$

Introducing the function of v_α

$$z[v_\alpha] := \left[\prod_{a=1}^n \int_0^{2\pi} \frac{d\lambda^a}{2\pi} e^{i\lambda^a} \right] \exp \left[v_\alpha e^{-i \sum_{b \in \alpha} \lambda^b} \right], \quad (32)$$

and the order parameters

$$Q_\alpha := \frac{X_\alpha + i Y_\alpha}{\sqrt{2}} \quad (33)$$

we can write

$$\begin{aligned} \overline{Z}^n &= \\ &= \left[\prod'_{\alpha \in \mathcal{P}([n])} \frac{N}{2\pi \hat{g}_\alpha} \iint dQ_\alpha dQ_\alpha^* \right] e^{-NS[Q] - N\Delta S^T[Q]}, \end{aligned} \quad (34a)$$

with

$$S[Q] = \sum_{\alpha \in \mathcal{P}([n])} \left(\frac{|Q_\alpha|^2}{\hat{g}_\alpha} - \ln z[Q_\alpha] - \ln z[Q_\alpha^*] \right) \quad (34b)$$

$$\Delta S^T[Q] = \sum'_{\substack{\alpha, \beta \in \mathcal{P}([n]) \\ \alpha \cap \beta = \emptyset}} \hat{g}_\alpha \hat{g}_\beta \frac{|Q_\alpha|^2}{4N \hat{g}_{\alpha \cup \beta}^2}. \quad (34c)$$

a form which is suitable to be evaluated, in the asymptotic limit for large N , by means of the saddle point method. It is immediately clear that ΔS^T contains a contribution to the action that is $O(N^{-1})$, and therefore it can be neglected in the evaluation of the leading contribution. It follows that the stationarity equations are of the form

$$\frac{Q_\alpha^*}{\hat{g}_\alpha} = \frac{d \ln z[Q_\alpha]}{d Q_\alpha}, \quad \frac{Q_\alpha}{\hat{g}_\alpha} = \frac{d \ln z[Q_\alpha^*]}{d Q_\alpha^*}. \quad (35)$$

The application of the saddle point method gives

$$\overline{Z}^n \simeq \exp \left(-NS[Q^{\text{sp}}] - N\Delta S^T[Q^{\text{sp}}] - \frac{1}{2} \ln \det \mathbf{\Omega}[Q^{\text{sp}}] \right), \quad (36)$$

where $\mathbf{\Omega}$ is the Hessian matrix of $S[Q]$, and Q^{sp} is the saddle point solution. As we will show below, the contribution $\ln \det \mathbf{\Omega}[Q^{\text{sp}}]$ provides finite size corrections to the leading contribution of the same order of the corrections in $N\Delta S^T[Q^{\text{sp}}]$.

A. Replica symmetric ansatz and limit of vanishing number of replicas

To proceed with our calculation, we adopt, as usual in the literature, a replica symmetric ansatz for the solution of the

saddle point equations. A replica symmetric solution is of the form

$$Q_\alpha = Q_\alpha^* = q_{|\alpha|}. \quad (37)$$

In particular this implies that $Y_\alpha = 0$. In order to analytically continue to $n \rightarrow 0$ the value at the saddle-point of S in Eq. (34b), let us first remark that under the assumption in Eq. (37)

$$\sum'_{\alpha \in \mathcal{P}([n])} \frac{|Q_\alpha|^2}{\hat{g}_\alpha} = \sum_{k=1}^n \binom{n}{k} \frac{q_k^2}{\hat{g}_k} = n \sum_{k=1}^{\infty} \frac{(-1)^{k-1} q_k^2}{k \hat{g}_k} + o(n). \quad (38)$$

Moreover, under our assumption (see Appendix A)

$$\sum'_{\alpha \in \mathcal{P}([n])} \ln z[Q_\alpha] = n \int_{-\infty}^{+\infty} (e^{-e^l} - e^{-G(l)}) dl, \quad (39)$$

where

$$G(l) := \sum_{k=1}^{\infty} (-1)^{k-1} q_k \frac{e^{lk}}{k!}. \quad (40)$$

In conclusion, under the replica symmetric ansatz in Eq. (37) the functional to be minimized is

$$\hat{\beta} \hat{F} = \sum_{k=1}^{\infty} \frac{(-1)^{k-1} q_k^2}{k \hat{g}_k} - 2 \int_{-\infty}^{+\infty} [e^{-e^l} - e^{-G(l)}] dl. \quad (41)$$

A variation with respect to q_k gives the saddle-point equation

$$\frac{1}{k} \frac{q_k}{\hat{g}_k} = \int_{-\infty}^{+\infty} e^{-G(y)} \frac{e^{yk}}{k!} dy \quad (42)$$

which is to say

$$\begin{aligned} G(l) &= \sum_{k=1}^{\infty} (-1)^{k-1} q_k \frac{e^{lk}}{k!} \\ &= \int_{-\infty}^{+\infty} e^{-G(y)} \sum_{k=1}^{\infty} (-1)^{k-1} k \hat{g}_k \frac{e^{(y+l)k}}{(k!)^2} dy. \end{aligned} \quad (43)$$

This implies that

$$\begin{aligned} \sum_{k=1}^{\infty} \frac{(-1)^{k-1} q_k^2}{k \hat{g}_k} &= \sum_{k=1}^{\infty} (-1)^{k-1} q_k \int_{-\infty}^{+\infty} e^{-G(y)} \frac{e^{yk}}{k!} dy \\ &= \int_{-\infty}^{+\infty} G(y) e^{-G(y)} dy. \end{aligned} \quad (44)$$

These formulas are for a general law ρ_r . Observe also that the expression of \hat{g}_p is not specified. For finite r and $N \rightarrow \infty$, Eq. (25) simplifies as

$$\lim_{N \rightarrow \infty} \hat{g}_p = \frac{\eta_0 \Gamma(r+1)}{(\hat{\beta} p)^{r+1}}. \quad (45)$$

We shall restrict to the case in which Eq. (45) holds. Then Eq. (43) becomes

$$G_r(l) = \frac{\eta_0 \Gamma(r+1)}{\hat{\beta}^{r+1}} \int_{-\infty}^{+\infty} B_r(l+y) e^{-G_r(y)} dy, \quad (46)$$

with

$$B_r(x) := \sum_{k=1}^{\infty} (-1)^{k-1} \frac{e^{xk}}{k^r (k!)^2}. \quad (47)$$

In Eq. (46), and in the following, we introduce the subindex r to stress the dependence of G and of the thermodynamical functionals on r . The average cost is therefore

$$\hat{E}_r = \frac{\partial}{\partial \hat{\beta}} \hat{\beta} \hat{F}_r = \frac{r+1}{\hat{\beta}} \int_{-\infty}^{+\infty} G_r(y) e^{-G_r(y)} dy. \quad (48)$$

Using the fact that (see Appendix B)

$$\lim_{\delta \rightarrow \infty} \frac{1}{\delta^r} B_r(\delta x) = \frac{x^r \theta(x)}{\Gamma(r+1)}, \quad (49)$$

if we introduce

$$\hat{G}_r(l) := G_r \left(\frac{\hat{\beta}}{[\eta_0 \Gamma(r+1)]^{\frac{1}{r+1}}} l \right) \quad (50)$$

in the limit $\hat{\beta} \rightarrow +\infty$ the function \hat{G}_r satisfies Eq. (12c) and the value of \hat{E}_r is the one reported in Eq. (12a). In particular, at fixed r , if we consider the two laws ρ_r^P and ρ_r^I , the ratio between the corresponding average optimal costs is given by

$$\lambda_r := \frac{\hat{E}_r^P}{\hat{E}_r^I} = \left(\frac{\eta_0^I}{\eta_0^P} \right)^{\frac{1}{r+1}} = [\Gamma(r+2)]^{-\frac{1}{r+1}}. \quad (51)$$

In the case $r = 0$, we have [8]

$$\hat{G}_0(l) = \ln(1 + e^l), \quad (52)$$

$$\hat{E}_0 = \frac{1}{\eta_0(0)} \int_{-\infty}^{+\infty} \frac{\ln(1 + e^y)}{1 + e^y} dy = \frac{1}{\eta_0(0)} \frac{\pi^2}{6}, \quad (53)$$

a result which has been obtained with a different method by Aldous [27]. For the evaluation of the integral see Appendix D.

III. FINITE SIZE CORRECTIONS

The evaluation of the first order corrections for finite number of points has been considered in Refs. [24, 25] in the $r = 0$ case. For this particular choice, and assuming a distribution law ρ_0^Γ , a much stronger conjecture was proposed by Parisi [28], and later proved by Linusson and Wästlund [29] and Nair *et al.* [30], that is, for every N

$$\hat{E}_0^\Gamma(N) = H_{N,2} := \sum_{k=1}^N \frac{1}{k^2}. \quad (54)$$

For large N , Parisi's formula implies

$$\hat{E}_0^\Gamma(N) = \frac{\pi^2}{6} - \frac{1}{N} + o\left(\frac{1}{N}\right). \quad (55)$$

Using instead the law ρ_0^P (uniform distribution on the interval) we have [24, 25]

$$\hat{E}_0^P(N) = \frac{\pi^2}{6} - \frac{1 + 2\zeta(3)}{N} + o\left(\frac{1}{N}\right) \quad (56)$$

from which we see that corrections for both laws are analytic, with the same inverse power of N , but of different coefficients.

In their study of the finite size corrections, the authors of Ref. [25] show that, in their particular case, there are two kind of finite size corrections. The first one comes from the application of the saddle point method, giving a series of corrections in the inverse powers of N . This contribution is the sum of two terms. The first term in this expansion corresponds to the contribution of the ΔS^T term given in Eq. (34c) appearing in the exponent in Eq. (36). The second term is related to the fluctuations, also appearing in Eq. (36), involving the Hessian of S . The second kind of corrections is due to the particular form of the law $\rho_r(w)$ for the random links, and in particular to the series expansion in Eq. (25). This contribution can be seen at the level of the action S in Eq. (34b), being

$$\begin{aligned} \frac{|Q_\alpha|^2}{\hat{g}_\alpha} &\approx |Q_\alpha|^2 \frac{(\hat{\beta}|\alpha|)^{r+1}}{\eta_0 \Gamma(r+1)} \\ &- |Q_\alpha|^2 \frac{(r+1)\eta_1}{N^{\frac{1}{r+1}} \eta_0^2} \frac{(\hat{\beta}|\alpha|)^r}{\Gamma(r+1)} + O\left(N^{-\frac{2}{r+1}}\right). \end{aligned} \quad (57)$$

In full generality, the expansion of $1/\hat{g}_\alpha$ generates a sum over terms each one of order $N^{-\frac{k}{r+1}}$ with $k \geq 1$. All these corrections are $o(N^{-1})$ for $r \in (-1, 0)$, whereas the corrections obtained from the contributions with $1 \leq k \leq r+1$ are of the same order of the analytic term, or greater, for $r \geq 0$. In particular, if $\eta_1 \neq 0$, for $r > 0$ the $k = 1$ term provides the leading correction, scaling as $N^{-\frac{1}{r+1}}$. It is also evident that all these corrections are absent if $\eta_k = 0$ for $k \geq 1$, as it happens in the case of the ρ_r^P law.

A. Correction due to η_1

Let us consider the $r \geq 0$ case and let us restrict ourselves to the $k = 1$ term, of order $N^{-\frac{1}{r+1}}$ in Eq. (13). Its contribution to the total free energy is given by

$$\hat{\beta} \Delta \hat{F}_r^{(1)} = -\frac{r+1}{N^{\frac{1}{r+1}}} \frac{\eta_1}{\eta_0^2} \sum_{p=1}^{\infty} \frac{(-1)^{p-1}}{p} \frac{(\hat{\beta}p)^r}{\Gamma(r+1)} q_p^2 \quad (58)$$

where we already made a replica symmetric assumption and considered the $n \rightarrow 0$ limit. Imposing the saddle point relation in Eq. (42), and using the limit in Eq. (45), we obtain

$$N^{\frac{1}{r+1}} \Delta \hat{F}_r^{(1)} =$$

$$\begin{aligned} &= -\frac{\eta_1(r+1)}{\hat{\beta}^2 \eta_0} \int_{-\infty}^{+\infty} e^{-G_r(y)} \sum_{p=1}^{\infty} \frac{(-1)^{p-1} q_p e^{py}}{p p!} dy \\ &= -\frac{\eta_1(r+1)}{\hat{\beta}^2 \eta_0} \int_{-\infty}^{+\infty} e^{-G_r(y)} \int_{-\infty}^y G_r(u) dy du \\ &= -\frac{\eta_1(r+1)}{\eta_0 [\eta_0 \Gamma(r+1)]^{\frac{2}{r+1}}} \int_{-\infty}^{+\infty} e^{-\hat{G}_r(y)} \int_{-\infty}^y \hat{G}_r(u) dy du. \end{aligned} \quad (59)$$

To put the expression above in the form presented in Eq. (13e), observe that

$$\begin{aligned} &\int_{-\infty}^{+\infty} e^{-\hat{G}_r(y)} \int_{-\infty}^y \hat{G}_r(u) dy du = \\ &= \int_{-\infty}^{+\infty} \hat{G}_r(-u) \int_{-\infty}^{+\infty} e^{-\hat{G}_r(y)} \theta(y+u) dy du \\ &= \int_{-\infty}^{+\infty} \hat{G}_r(-u) \mathcal{D}_u^r \hat{G}_r(u) du \equiv J_r^{(r)}, \end{aligned} \quad (60)$$

a structure that can be more useful for numerical evaluation, at least for r integer. In the previous equation we have used Eq. (C4) and we have introduced the Riemann–Liouville fractional derivative

$$\begin{aligned} \mathcal{D}_t^\alpha f(t) &:= \frac{d^{[\alpha]+1}}{dt^{[\alpha]+1}} \int_{-\infty}^t \frac{(t-\tau)^{[\alpha]-\alpha}}{\Gamma([\alpha]-\alpha+1)} f(\tau) d\tau, \\ \alpha &\geq 0, \quad f \in L_p(\Omega) \quad \forall p \in \left[1, \frac{1}{[\alpha]-\alpha+1}\right), \end{aligned} \quad (61)$$

where $\Omega := (-\infty, t)$ is the domain of integration (see the monographs in Refs. [31, 32] for further details).

B. Correction due to the saddle point approximation

Let us now consider the corrections due to the saddle point approximation. The first contribution is expressed by ΔS^T , given in Eq. (34c). In the replica symmetric hypothesis, we have that

$$\begin{aligned} &\sum'_{\substack{\alpha, \beta \in \mathcal{P}(\{n\}) \\ \alpha \cap \beta = \emptyset}} \hat{g}_\alpha \hat{g}_\beta \frac{|Q_\alpha|^2}{4N \hat{g}_{\alpha \cup \beta}^2} = \\ &= \frac{1}{N} \sum_{s=1}^{\infty} \sum_{t=1}^{\infty} \binom{n}{s, t, n-s-t} \frac{\hat{g}_s \hat{g}_t}{\hat{g}_{s+t}^2} q_{s+t}^2. \end{aligned} \quad (62)$$

We can write the corresponding correction to the free energy as

$$\Delta \hat{F}_r^T = \frac{1}{2\hat{\beta}N} \sum_{s=1}^{\infty} \sum_{t=1}^{\infty} (-1)^{s+t-1} \frac{(s+t-1)!}{s! t!} \frac{\hat{g}_s \hat{g}_t}{\hat{g}_{s+t}^2} q_{s+t}^2. \quad (63)$$

In Appendix E we show that the previous quantity can be written as

$$\begin{aligned} \Delta \hat{F}_r^T &= -\frac{\Gamma(2r+2)}{N\eta_0^{\frac{1}{r+1}}\Gamma^{2+\frac{1}{r+1}}(r+1)} \frac{1}{r+1} \int_{-\infty}^{+\infty} \hat{G}_r(-u)\hat{G}_r(u) du \\ &= -\frac{\Gamma(2r+2)}{N\eta_0^{\frac{1}{r+1}}\Gamma^{2+\frac{1}{r+1}}(r+1)} \frac{J_r^{(0)}}{r+1}. \end{aligned} \quad (64)$$

Another type of finite size correction comes from the fluctuations around the saddle point [25, Section B.3], related to the Hessian matrix $\hat{\Omega}$ appearing in Eq. (36). The evaluation of the contribution of the Hessian matrix is not trivial and it has been discussed by Mézard and Parisi [24] and, later, by Parisi and Ratiéville [25]. They proved that the whole contribution comes from a volume factor due to a non-trivial metric $\hat{\Omega}$ obtained from Ω imposing the replica symmetric assumption and such that

$$\ln \sqrt{\det \hat{\Omega}} = \ln \sqrt{\det \Omega}. \quad (65)$$

The $n \times n$ matrix $\hat{\Omega}$ can be written as

$$\hat{\Omega} = na_1 \mathbf{\Pi} + (a_0 - a_1) \mathbb{I}_n, \quad (66)$$

where \mathbb{I}_n is the $n \times n$ identity matrix, we have introduced the quantities

$$a_0 := \sum_{p=1}^{\infty} \binom{n-1}{p-1} \frac{q_p^2}{\hat{g}_p} \quad (67a)$$

$$a_1 := \sum_{p=2}^{\infty} \binom{n-2}{p-2} \frac{q_p^2}{\hat{g}_p}, \quad (67b)$$

and $\mathbf{\Pi}$ is a projection matrix on the constant vector defined as

$$\mathbf{\Pi} := \frac{\mathbb{J}_n}{n}, \quad (68)$$

where \mathbb{J}_n is the $n \times n$ matrix with all entries equal to 1. The matrix $\mathbf{\Pi}$ has 1 eigenvalue equal to 1 and $n-1$ eigenvalues equal to 0. It follows that, because the two matrices $\mathbf{\Pi}$ and \mathbb{I}_n obviously commute, $\hat{\Omega}$ has 1 eigenvalue equal to $a_0 + (n-1)a_1$ and $n-1$ eigenvalues equal to $a_0 - a_1$. Its determinant is therefore simply given by

$$\det \hat{\Omega} = (a_0 - a_1)^{n-1} [a_0 + (n-1)a_1]. \quad (69)$$

In the limit of $n \rightarrow 0$ we easily get

$$\begin{aligned} a_0 &= \sum_{p=1}^{\infty} (-1)^{p-1} \frac{q_p^2}{\hat{g}_p} = \sum_{p=1}^{\infty} (-1)^{p-1} p q_p \int_{-\infty}^{+\infty} e^{-G_r(y)} \frac{e^{py}}{p!} dy \\ &= \int_{-\infty}^{+\infty} e^{-G_r(y)} \frac{dG_r(y)}{dy} dy = \int_{-\infty}^{+\infty} e^{-\hat{G}_r(y)} \frac{d\hat{G}_r(y)}{dy} dy \\ &= - \int_{-\infty}^{+\infty} \frac{d}{dy} e^{-\hat{G}_r(y)} dy = e^{-\hat{G}_r(-\infty)} - e^{-\hat{G}_r(+\infty)} = 1 \end{aligned} \quad (70)$$

for all values of r . Similarly,

$$a_1 = - \sum_{p=2}^{\infty} (-1)^{p-1} (p-1) \frac{q_p^2}{\hat{g}_p} = - \sum_{p=1}^{\infty} (-1)^{p-1} (p-1) \frac{q_p^2}{\hat{g}_p} \quad (71)$$

so that

$$\begin{aligned} a_0 - a_1 &= \sum_{p=1}^{\infty} (-1)^{p-1} p \frac{q_p^2}{\hat{g}_p} \\ &= \sum_{p=1}^{\infty} (-1)^{p-1} p^2 q_p \int_{-\infty}^{+\infty} e^{-G_r(y)} \frac{e^{py}}{p!} dy \\ &= \int_{-\infty}^{+\infty} e^{-G_r(y)} \frac{d^2}{dy^2} G_r(y) dy \\ &= \frac{[\eta_0 \Gamma(r+1)]^{\frac{1}{r+1}}}{\hat{\beta}} \int_{-\infty}^{+\infty} e^{-\hat{G}_r(y)} \frac{d^2}{dy^2} \hat{G}_r(y) dy. \end{aligned} \quad (72)$$

Therefore

$$\sqrt{\det \hat{\Omega}} = 1 + \frac{n}{2} \left[\frac{a_1}{a_0 - a_1} + \ln(a_0 - a_1) \right] + o(n). \quad (73)$$

In conclusion, integrating by parts and using Eq. (C6), we obtain

$$\begin{aligned} \Delta \hat{F}_r^F &= - \lim_{\hat{\beta} \rightarrow \infty} \frac{1}{nN\hat{\beta}} \ln \sqrt{\det \hat{\Omega}} \\ &= - \frac{1}{2N [\eta_0 \Gamma(r+1)]^{\frac{1}{r+1}} J_r^{(r+3)}}. \end{aligned} \quad (74)$$

C. Application: the $r = 0$ case

The results obtained in the $r = 0$ case, analyzed by Parisi and Ratiéville [25], can be easily recovered. From the general expression in Eq. (13), by setting $r = 0$, we get

$$\begin{aligned} \Delta \hat{F}_0 &:= \Delta \hat{F}_0^{(1)} + \Delta \hat{F}_0^T + \Delta \hat{F}_0^F \\ &= - \frac{1}{\eta_0(0)N} \left[\left(1 + \frac{\eta_1(0)}{\eta_0(0)} \right) \frac{J_0^{(0)}}{\eta_0(0)} + \frac{1}{2J_0^{(3)}} \right] \\ &= - \frac{1}{\eta_0(0)N} \left[\left(1 + \frac{\eta_1(0)}{\eta_0(0)} \right) \frac{2\zeta(3)}{\eta_0(0)} + 1 \right], \end{aligned} \quad (75)$$

where we have used the results discussed in the Appendix D for the two integrals involved in the expression above. Eq. (55) and Eq. (56) are obtained using Eq. (16) and Eq. (17) respectively.

IV. THE LIMITING CASE $r \rightarrow +\infty$

In this section we shall concentrate to the limiting case in which $r \rightarrow +\infty$. We can easily verify that, in the weak sense,

$$\lim_{r \rightarrow +\infty} \rho_r^P(w) = \delta(w-1) \quad (76)$$

so that all the weights become equal to unity. We expect therefore that

$$\lim_{r \rightarrow +\infty} \hat{E}_r^{\text{P}}(N) = 1, \quad (77)$$

independently from N . The average cost obtained using ρ_r^{Γ} , instead, diverges, and it is therefore more interesting to consider the modified law

$$\rho_r^{\gamma}(w) := \frac{(r+1)^{r+1}}{\Gamma(r+1)} w^r e^{-(r+1)w} \theta(w) \xrightarrow{r \rightarrow \infty} \delta(w-1). \quad (78)$$

According to our general discussion, we have that

$$\eta_k^{\gamma}(r) = \frac{(r+1)^{k+r+1}}{\Gamma(r+1)} \frac{(-1)^k}{k!}, \quad k \geq 0, \quad (79)$$

implying that, independently from N ,

$$\hat{E}_r^{\gamma}(N) = \frac{1}{r+1} \hat{E}_r^{\Gamma}(N), \quad (80)$$

and therefore

$$\hat{E}_r^{\gamma} = \frac{\Gamma(r+2)^{\frac{1}{r+1}}}{r+1} \hat{E}_r^{\text{P}} \quad (81)$$

in the limit of infinite N . In particular

$$\lim_{r \rightarrow +\infty} \hat{E}_r^{\gamma} = \frac{1}{e}. \quad (82)$$

It follows that, even though the two laws ρ^{P} and ρ^{γ} both converge to the same limiting distribution, according to our formulas the corresponding average cost is *not* the same. This is due to the fact that the two limits $N \rightarrow +\infty$ and $r \rightarrow +\infty$ do not commute for the law ρ^{γ} , because of the presence of $O(N^{-\frac{k}{r+1}})$ corrections that give a leading contribution if the $r \rightarrow \infty$ limit is taken first.

To look into more details in the $r \rightarrow +\infty$ limit, we find convenient, when looking at the saddle point solution, to perform a change of variables, following the approach in Refs. [33, 34], that is to write

$$\mathcal{G}_r(x) := \hat{G}_r \left[\Gamma^{\frac{1}{r+1}}(r+2) \left(\frac{1}{2} + \frac{x}{r+1} \right) \right] \quad (83)$$

then Eq. (12c) becomes

$$\mathcal{G}_r(x) = \int_{-x-r-1}^{+\infty} \left(1 + \frac{x+t}{r+1} \right)^r e^{-\mathcal{G}_r(t)} dt, \quad (84)$$

so that, in the $r \rightarrow +\infty$ limit

$$\mathcal{G}_{\infty}(x) = e^x \int_{-\infty}^{+\infty} e^{t-\mathcal{G}_{\infty}(t)} dt. \quad (85)$$

If we set $\mathcal{G}_{\infty}(x) = ae^x$, with

$$a = \int_{-\infty}^{+\infty} e^{t-\mathcal{G}_{\infty}(t)} dt \quad (86)$$

we recover

$$a = \int_{-\infty}^{+\infty} e^{t-ae^t} dt = \int_0^{+\infty} e^{-az} dz = \frac{1}{a} \Rightarrow \mathcal{G}_{\infty}(x) = e^x. \quad (87)$$

From Eq. (12a) with the change of variable in Eq. (83), we get

$$\hat{E}_r = \left(\frac{r+1}{\eta_0(r)} \right)^{\frac{1}{r+1}} \int_{-\infty}^{+\infty} \mathcal{G}_r(x) e^{-\mathcal{G}_r(x)} dx, \quad (88)$$

so that

$$\hat{E}_{\infty} = \lim_{r \rightarrow +\infty} \left(\frac{r+1}{\eta_0(r)} \right)^{\frac{1}{r+1}} = \lim_{r \rightarrow +\infty} \eta_0(r)^{-\frac{1}{r+1}} = \begin{cases} 1 & \text{for } \rho_r^{\text{P}}, \\ \frac{1}{e} & \text{for } \rho_r^{\gamma}, \end{cases} \quad (89)$$

in agreement with the previous results.

Let us now evaluate the integrals appearing in the finite size corrections in Eq. (13). Let us first start with the ΔF_r^T and the ΔF_r^F corrections. From the definition, for large r

$$\begin{aligned} J_r^{(0)} &= \int_{-\infty}^{+\infty} \hat{G}_r(y) \hat{G}_r(-y) dy \\ &= \int_{-\infty}^{+\infty} dy \int_{-y}^{\infty} dt_1 \frac{(t_1+y)^r}{\Gamma(r+1)} e^{-\hat{G}_r(t_1)} \int_y^{\infty} dt_2 \frac{(t_2+y)^r}{\Gamma(r+1)} e^{-\hat{G}_r(t_2)} \\ &= \Gamma^{\frac{1}{r+1}}(r+2) \int_{-\infty}^{+\infty} dt_1 \int_{-\infty}^{+\infty} dt_2 K_r(t_1, t_2) e^{-\mathcal{G}_r(t_1) - \mathcal{G}_r(t_2)} \quad (90) \end{aligned}$$

where

$$\begin{aligned} K_r(t_1, t_2) &:= \int_{-x_1-r-1}^{x_2} \frac{(t+t_1+r+1)^r (t_2-t)^r}{(r+1)^{2r+1}} dt \\ &= \left(1 + \frac{x_1+x_2}{r+1} \right)^{2r+1} \frac{\Gamma^2(r+1)}{\Gamma(2r+2)} \quad (91) \end{aligned}$$

and therefore for large r

$$\begin{aligned} J_r^{(0)} &\simeq \Gamma^{\frac{1}{r+1}}(r+2) \frac{\Gamma^2(r+1)}{\Gamma(2r+2)} \left(\int_{-\infty}^{+\infty} e^{x-e^x} dx \right)^2 \\ &= \Gamma^{\frac{1}{r+1}}(r+2) \frac{\Gamma^2(r+1)}{\Gamma(2r+2)} \simeq \frac{\Gamma^{2+\frac{1}{r+1}}(r+1)}{\Gamma(2r+2)}, \quad (92) \end{aligned}$$

so we get

$$\Delta \hat{F}_r^T \simeq -\frac{1}{N \eta_0^{\frac{1}{r+1}}(r)} \quad (93)$$

a contribution which vanishes as r^{-1} both for the law ρ_r^{P} and the law ρ_r^{γ} (indeed we know that, in this case, all corrections

must vanish when $r \rightarrow \infty$ at fixed N). In particular, if we consider the law ρ_r^Γ , we have, in the $r \rightarrow \infty$ limit,

$$\Delta \hat{F}_\infty^T = -\frac{1}{eN}. \quad (94)$$

Similarly, for large r , we have that

$$\frac{1}{J_r^{(r+3)}} = \frac{\Gamma(r+2)^{\frac{1}{r+1}}}{r+1} \left[\int_{-\infty}^{+\infty} e^{-\mathcal{G}_r(x)} \left(\frac{d}{dx} \right)^2 \mathcal{G}_r(x) \right]^{-1} = \frac{1}{e} \quad (95)$$

and therefore

$$\Delta \hat{F}_r^F \simeq -\frac{1}{2N\eta_0^{\frac{1}{r+1}}(r)r} = \frac{\Delta \hat{F}_r^T}{2}. \quad (96)$$

Instead, if we consider $\Delta F_r^{(1)}$, we have that

$$J_r^{(r)} = \left[\frac{\Gamma(r+2)^{\frac{1}{r+1}}}{r+1} \right]^2 \int_{-\infty}^{+\infty} du e^{-\mathcal{G}_r(u)} \int_{-\infty}^u dv \mathcal{G}_r(v) \xrightarrow{r \rightarrow \infty} \frac{1}{e^2} \int_{-\infty}^{+\infty} du e^{-e^u} \int_{-\infty}^u dv e^v = \frac{1}{e^2}, \quad (97)$$

finally obtaining

$$\Delta \hat{F}_r^{(1)} \simeq -\frac{\eta_1(r)}{N^{\frac{1}{r+1}} \eta_0^{1+\frac{2}{r+1}}(r)r} \quad (98)$$

so that, considering the law ρ_r^γ , if we send $r \rightarrow \infty$ before taking the limit $N \rightarrow \infty$, $\Delta \hat{F}_\infty^{(1)} \sim O(1)$ and we get a new contribution to the average optimal cost

$$\hat{E} = \hat{E}_\infty + \sum_{k=1}^{\infty} \Delta \hat{F}_\infty^{(k)} = \frac{1}{e} + \frac{1}{e^2} + \dots \quad (99)$$

a series where we miss the contributions of order $N^{\frac{k}{r+1}}$ for $k \geq 2$, and that we know will sum up to 1.

V. NUMERICAL RESULTS

In this Section we discuss some numerical results. First, we present a numerical study of our theoretical predictions obtained in the previous Sections. Second, we compare with numerical simulations, in which the random assignment problem is solved using an exact algorithm.

The evaluation of all quantities in Eq. (13) depends on the solution of Eq. (12c). We solved numerically this equation for general r by a simple iterative procedure. In particular, for $r > 0$ we generated a grid of $2K - 1$ equispaced points in an interval $[-y_{\max}, y_{\max}]$ and we used a discretized version of the saddle point equation in Eq. (12c) in the form

$$\hat{G}_r^{[s+1]}(y_i) = \frac{y_{\max}}{K} \sum_{k=2K-i}^{2K} \frac{(y_i + y_k)^r e^{-\hat{G}_r^{[s]}(y_k)}}{\Gamma(r+1)} \quad (100)$$

$$y_i = \frac{i-K}{K} y_{\max}, \quad i = 0, 1, \dots, 2K.$$

We imposed as initial function $\hat{G}_r^{[0]}$ of the iterative procedure

$$\hat{G}_r^{[0]}(y_i) \equiv \hat{G}_0(y_i) = \ln(1 + e^{y_i}). \quad (101)$$

We observed that the quantity

$$\Delta \hat{G}_r^{[s]} = \sum_{i=0}^{2K} \left| \hat{G}_r^{[s]}(y_i) - \hat{G}_r^{[s-1]}(y_i) \right| \quad (102)$$

decays exponentially with s , and therefore convergence is very fast. For our computation, we used typically 30 iteration.

For $r < 0$ the term $(l+y)^r$ in the saddle point equation is divergent in $y = -l$ and Eq. (100) cannot be adopted. We have therefore rewritten the saddle point equation using an integration by parts, obtaining

$$\hat{G}_r(l) = \int_{-l}^{+\infty} \frac{(l+y)^{r+1} e^{-\hat{G}_r(y)} d\hat{G}_r(y)}{\Gamma(r+2)} \frac{d\hat{G}_r(y)}{dy} dy. \quad (103)$$

After discretizing the previous equation, we used the same algorithm described for the $r \geq 0$ case (for a discussion on the uniqueness of the solution of Eq. (12c), see Ref. [35]).

In Table I we present our numerical results for the quantities

$$\eta_0^{\frac{1}{r+1}} \hat{E}_r, \quad (N\eta_0^2)^{\frac{1}{r+1}} \frac{\eta_0 \Delta \hat{F}_r^{(1)}}{\eta_1}, \quad N\eta_0^{\frac{1}{r+1}} \Delta \hat{F}_r^T, \quad N\eta_0^{\frac{1}{r+1}} \Delta \hat{F}_r^F$$

for different values of r . Observe that the quantities appearing in the expansion in Eqs. (13) can be calculated using these values for any ρ_r at given r , in addition to simple prefactors depending on the chosen distribution ρ_r .

In order to test our analysis for correction terms, we performed direct Monte Carlo sampling on a set of instances. Previous simulations have been reported, for example, in Refs. [8, 33, 36, 37]. In our setting, each realization of the matching problem has been solved by a C++ implementation of the Jonker-Volgenant algorithm [38].

We first evaluated the asymptotic average optimal costs \hat{E}_r^P and \hat{E}_r^Γ , obtained, for different values of r , using the laws ρ_r^P and ρ_r^Γ respectively. In the case of the law ρ_r^P , the asymptotic estimate for \hat{E}_r^P has been obtained using the fitting function

$$f^P(N) = \alpha_r^P + \frac{\beta_r^P}{N}, \quad (104)$$

with α_r^P and β_r^P fitting parameters to be determined, α_r^P corresponding to the value of the average optimal cost in the $N \rightarrow \infty$ limit. For a given value of r , we averaged over I_N instances for each value of N accordingly with the table below.

N	500	750	1000	2500	5000
I_N	100000	75000	50000	20000	10000

r	$\eta_0^{\frac{1}{r+1}} \hat{E}_r$	$(N\eta_0^2)^{\frac{1}{r+1}} \frac{\eta_0 \Delta \hat{E}_r^{(1)}}{\eta_1}$	$N\eta_0^{\frac{1}{r+1}} \Delta \hat{F}_r^T$	$N\eta_0^{\frac{1}{r+1}} \Delta \hat{F}_r^F$
-0.5	1.125775489	-2.777285153	-3.917446075	-1.192663973
-0.4	1.334614017	-2.952484269	-3.665262242	-1.250475151
-0.3	1.471169704	-2.921791666	-3.324960744	-1.222990786
-0.2	1.558280634	-2.784084499	-2.984917100	-1.157857158
-0.1	1.612502443	-2.600804197	-2.675513663	-1.079610016
0	1.644934067	-2.404113806	-2.404113806	-1
0.1	1.662818967	-2.215821874	-2.168528577	-0.924257491
0.2	1.671039856	-2.038915744	-1.966713438	-0.854501434
0.3	1.672729262	-1.877696614	-1.792481703	-0.791231720
0.4	1.670005231	-1.732453452	-1.641566768	-0.734262435
0.5	1.664311154	-1.602337915	-1.510248399	-0.683113178
0.6	1.656639222	-1.486024319	-1.395391897	-0.637204338
0.7	1.647677145	-1.382051819	-1.294397704	-0.595951390
0.8	1.637905005	-1.288993419	-1.205124002	-0.558807473
0.9	1.627659755	-1.205532353	-1.125808312	-0.525279810
1	1.617178636	-1.130489992	-1.054997763	-0.494933215
2	1.519733739	-0.670341811	-0.626403698	-0.303146650
3	1.446919560	-0.461144035	-0.431759755	-0.211631545
4	1.393163419	-0.346056113	-0.324185048	-0.159938240
5	1.352087648	-0.274505368	-0.257174804	-0.127356338
6	1.319651066	-0.226200326	-0.211931870	-0.105200594
7	1.293333076	-0.191617643	-0.179566694	-0.089276830
8	1.271505390	-0.165752490	-0.155385461	-0.077340947
9	1.253073980	-0.145742887	-0.136697943	-0.068095120
10	1.237277174	-0.129842072	-0.121861122	-0.060741591

Table I. Numerical values of the rescaled corrections appearing in Eqs. (13) for different values of r .

Similarly, the asymptotic average optimal cost \hat{E}_r^Γ has been obtained using a fitting function in the form

$$f^\Gamma(N) = \begin{cases} \alpha_r^\Gamma + \beta_r^\Gamma N^{-1} + \gamma_r^\Gamma N^{-\frac{1}{r+1}} & \text{for } -\frac{1}{2} \leq r < 1, \\ \alpha_r^\Gamma + \gamma_r^\Gamma N^{-\frac{1}{r+1}} + \delta_r^\Gamma N^{-\frac{2}{r+1}} & \text{for } r \geq 1. \end{cases} \quad (105)$$

We adopted therefore a three parameters fitting function, constructed according to Eq. (13) including the finite size correction up to order $o(N^{-1})$ for $r \geq 0$ and up to order $O(N^{-2})$ for $\frac{1}{2} \leq r < 2$. As in the case before, the asymptotic estimation for \hat{E}_r^Γ is given by α_r^Γ . Our data were obtained extrapolating the $N \rightarrow \infty$ limit from the average optimal cost for different values of N . The investigated sizes and the number of iterations were the same adopted for the evaluation of \hat{E}_r^P . To better exemplify the main differences in the finite size scaling between the ρ_r^P -case and the ρ_r^Γ -case, we have presented the numerical results for $r = 1$ in Fig. 1. In the picture, it is clear that the asymptotic value $\hat{E}_1^P = \frac{1}{\sqrt{2}} \hat{E}_1^\Gamma$ is the same in the two cases, as expected by Eq. (51), but the finite size corrections are different both in sign and in their scaling properties. In Table II we compare the results of our numerical simulations with the ones in the literature (when available) for both $\hat{E}_r^P(N)$ and $\lambda_r \hat{E}_r^\Gamma(N)$, λ_r being defined in Eq. (51). In Fig. 2 we plot our theoretical predictions and the numerical results presented in Table II.

Let us now consider the finite size corrections. In the case of the ρ_r^P law, the $O(N^{-1})$ corrections are given by $\Delta \hat{F}_r^T + \Delta \hat{F}_r^F$




Figure 1. Numerical results for $\hat{E}_1^P(N)$ and $\hat{E}_1^\Gamma(N)$ for several values of N . Note that finite size corrections have different sign for $N \rightarrow +\infty$. In the picture, we have represented also the theoretical predictions for both cases obtained including the finite size corrections up to $O(N^{-1})$.




Figure 2. Theoretical prediction of \hat{E}_r^P for several values of r (black line), compared with our numerical results. The dashed line is the large r asymptotic estimate, equal to 1. Error bars do not appear because smaller than the marks in the plot. The values for $\lambda_r \hat{E}_r^\Gamma$ almost coincide with the values of \hat{E}_r^P (see Table II) and are not represented.

and no nonanalytic corrections to the leading term appear. We obtain the finite size corrections from the data used for Table II, using Eq. (104) but fixing α_r^P to the average optimal cost \hat{E}_r^P given by the theoretical prediction in Table I and therefore with one free parameter only, namely β_r^P . In Fig. 3 we compare our predictions for $\Delta \hat{F}_r^T + \Delta \hat{F}_r^F$, deduced by the values in Table I, with the results of our numerical simulations for different values of r .

In the case of the ρ_r^Γ law with $r > 0$, the first correction to the average optimal cost is given by $\Delta \hat{F}_r^{(1)}$, whereas $\Delta \hat{F}_r^{(1)}$

r	\hat{E}_r^P	$\lambda_r \hat{E}_r^\Gamma$	\hat{E}_r^P [33]	\hat{E}_r^P [37]	Th. prediction
-0.5	4.5011(3)	4.504(1)	–	–	4.503101957
-0.4	3.12611(5)	3.1268(2)	–	–	3.126825159
-0.3	2.4484(1)	2.4488(3)	–	–	2.448788557
-0.2	2.0593(5)	2.0593(3)	–	–	2.059601452
-0.1	1.8127(3)	1.8126(2)	–	–	1.812767212
0	1.64500(5)	1.6449(2)	1.645(1)	1.6450(1)	1.644934067
0.1	1.5245(2)	1.5253(9)	–	–	1.524808331
0.2	1.4356(2)	1.4357(5)	–	–	1.435497487
0.3	1.3670(1)	1.3670(4)	–	–	1.367026464
0.4	1.31323(6)	1.3132(3)	–	–	1.3132296
0.5	1.27007(8)	1.2697(4)	–	–	1.270107121
0.6	1.2350(1)	1.2348(3)	–	–	1.234960167
0.7	1.20585(6)	1.2062(6)	–	–	1.205907312
0.8	1.18143(3)	1.1812(7)	–	–	1.181600461
0.9	1.16099(8)	1.1605(6)	–	–	1.161050751
1	1.14344(7)	1.1433(4)	1.143(2)	–	1.14351798
2	1.05371(1)	1.054(1)	1.054(1)	1.054(1)	1.053724521
3	1.02311(1)	1.0288(9)	1.0232(1)	1.0236(2)	1.023126632
4	1.009690(4)	1.010(3)	1.0098(1)	–	1.009736514
5	1.00303(2)	1.005(3)	1.00306(8)	1.0026(8)	1.003027802

Table II. Numerical results for the average optimal cost for different values of r , and theoretical predictions. The value \hat{E}_r^P from Ref. [33], due to a different convention adopted in their paper, is obtained as

$$\hat{E}_r^P = \left(\frac{2\pi^{\frac{r+1}{2}} \Gamma(r+1)}{\Gamma(\frac{r+1}{2}) \Gamma(r+2)} \right)^{\frac{1}{r+1}} \beta_{\text{num}}(r+1)$$

from Table II therein. The data for \hat{E}_r^P from Ref. [37] have been obtained via a linear fit, using a fitting function in the form of Eq. (104).

is $o(N^{-1})$ for $r < 0$. Again, this correction can be obtained by a fit of the same data used to extrapolate the average optimal cost, fixing the fitting parameter α_r^Γ in Eq. (105) to the theoretical prediction, and performing a two parameters fit in which the quantity γ_r^Γ appearing in Eq. (105) corresponds to $\Delta \hat{F}_r^{(1)}$. In Fig. 4 we compare our prediction for $\Delta \hat{F}_r^{(1)}$, given in Table I, with the results of our fit procedure for γ_r^Γ for $-\frac{1}{2} < r \leq 5$. Observe that the numerical evaluation of the single contribution $\Delta \hat{F}_r^{(1)}$ is not possible for $r = 0$. In this case, the result of our fit for the $O(N^{-1})$ correction was $\beta_r^\Gamma + \gamma_r^\Gamma = -0.97(4)$, to be compared with the theoretical prediction $N(\Delta \hat{F}_0^{(1)} + \Delta \hat{F}_0^F + \Delta \hat{F}_0^T) = -0.998354732 \dots$

VI. CONCLUSIONS

In the present paper we have discussed the finite size corrections in the random assignment problem for a generic distribution law $\rho_r(w)$ for the weights in the form of Eq. (11). We have shown that, for $r > 0$ and $\eta_1 \neq 0$, the first finite size correction scales as $O(N^{-\frac{1}{r+1}})$ and it is proportional to η_1 . In particular, the integrals $J_r^{(0)}$, $J_r^{(r)}$ and $J_r^{(r+3)}$ are positive quantities (see Appendix C). Therefore, independently from r , the corrections $\Delta \hat{F}_r^T$ and $\Delta \hat{F}_r^F$ are negative, while $\Delta \hat{F}_r^{(1)}$ has opposite sign respect to η_1 , so that, for example, it is positive for the law ρ_r^Γ , while it vanishes for the law ρ_r^P . We




Figure 3. Numerical estimates of $\Delta \hat{F}_r^T + \Delta \hat{F}_r^F$ for several values of r (red line), and theoretical prediction (blue line) obtained using the law ρ_r^P . The dashed line is the large r asymptotic estimate.




Figure 4. Numerical estimates of $\Delta \hat{F}_r^{(1)}$ for several values of r (red), and theoretical prediction (blue) obtained using the law ρ_r^Γ . Observe that a discrepancy between the theoretical prediction and the numerical results appear for $r \geq 1$: we interpret this fact as a consequence of the similar scaling of $\Delta \hat{F}_r^{(1)}$ and $\Delta \hat{F}_r^{(2)}$ for $r \gg 1$, that makes the numerical evaluation of the single contribution $\Delta \hat{F}_r^{(1)}$ difficult. The dashed line is the large r asymptotic estimate.

also provided a general expression for the coefficients of the $O(N^{-\frac{1}{r+1}})$ and $O(N^{-1})$ corrections. Moreover, we have shown that, if $\lim_{r \rightarrow +\infty} \rho_r = \rho$, then in general

$$\lim_{r \rightarrow \infty} \lim_{N \rightarrow \infty} \hat{E}_r(N) \neq \lim_{N \rightarrow \infty} \lim_{r \rightarrow \infty} \hat{E}_r(N). \quad (106)$$

We have finally numerically verified our results, by a numerical integration of our formulas and a comparison with simulations.

The $O(N^{-\frac{k}{r+1}})$ corrections appearing in Eq. (13), for $2 \leq k \leq [r] + 1$, remain to be computed. As discussed above, in the

$r \rightarrow \infty$ limit, it is expected that all these finite size corrections contribute to the leading term, justifying the noncommutativity of the limits in Eq. (106).

VII. ACKNOWLEDGMENTS

The authors thank Giorgio Parisi for many discussions and suggestions. The authors also thank Carlo Lucibello and Marco Gherardi for discussions on the numerical solution of the integral equations involving \hat{G}_r . M.P.D. thanks the LCM for the extensive use of their computational resources. The work of G.S. was supported by a grant from the Simons Foundation (#454949, Giorgio Parisi).

Appendix A: Evaluation of $z[Q]$ on the saddle point and analytic continuation for $n \rightarrow 0$

Let us evaluate now the quantity $z[Q]$ on the saddle point. Using the fact that, for any analytic function f

$$\int_0^{2\pi} \frac{d\lambda}{2\pi} e^{i\lambda} f(e^{-i\lambda}) = \oint \frac{d\xi}{2\pi i} \frac{f(\xi)}{\xi^2} = \left. \frac{df}{d\xi} \right|_{\xi=0} \quad (\text{A1})$$

we can write

$$\begin{aligned} & \left[\prod_{a=1}^n \int_0^{2\pi} \frac{d\lambda^a}{2\pi} e^{i\lambda^a} \right] \exp \left\{ \sum'_{\alpha \in \mathcal{P}([n])} q_{|\alpha|} e^{-i \sum_{b \in \alpha} \lambda^b} \right\} \\ &= \left. \frac{\partial^n}{\partial \xi_1 \cdots \partial \xi_n} \right|_{\xi_1 = \cdots = \xi_n = 0} \exp \left\{ \sum'_{\alpha \in \mathcal{P}([n])} q_{|\alpha|} \prod_{b \in \alpha} \xi_b \right\} \\ &= \sum_{\alpha} \prod_{\alpha_i \in \alpha} q_{|\alpha_i|}, \quad (\text{A2}) \end{aligned}$$

where $\alpha = \{\alpha_i\}_i$, $\alpha_i \in \mathcal{P}([n])$ are disjoint subsets whose union is $[n]$. But

$$\begin{aligned} \sum_{\alpha} \prod_{\alpha_i \in \alpha} q_{|\alpha_i|} &= \sum_{m=1}^n \sum_{\substack{k_1, \dots, k_m \\ k_1 + \dots + k_m = n}} \binom{n}{k_1 \dots k_m} \frac{q_{k_1} \cdots q_{k_m}}{m!} \\ &= \left(\frac{d}{dt} \right)^n \sum_{m=0}^{\infty} \frac{1}{m!} \sum_{k_1, \dots, k_m} \frac{q_{k_1} \cdots q_{k_m} t^{k_1 + \dots + k_m}}{k_1! \cdots k_m!} \Bigg|_{t=0} \\ &= \left(\frac{d}{dt} \right)^n \sum_{m=0}^{\infty} \frac{1}{m!} \left(\sum_{k=1}^{\infty} q_k \frac{t^k}{k!} \right)^m \Bigg|_{t=0} \\ &= \left(\frac{d}{dt} \right)^n \exp \left(\sum_{k=1}^{\infty} q_k \frac{t^k}{k!} \right) \Bigg|_{t=0}. \quad (\text{A3}) \end{aligned}$$

To perform the analytic prolongation, we prove now that, if $f(0) = 1$, then

$$\lim_{n \rightarrow 0} \frac{1}{n} \ln \left[\left(\frac{d}{dt} \right)^n f(t) \Bigg|_{t=0} \right] = \int_{-\infty}^{+\infty} [e^{-e^l} - f(-e^l)] dl. \quad (\text{A4})$$

This fact can be seen observing that, for $n \rightarrow 0$,

$$\begin{aligned} \left(\frac{d}{dt} \right)^n f(t) &= f \left(\frac{\partial}{\partial J} \right) J^n e^{Jt} \Bigg|_{J=0} \\ &\approx f(t) + n f \left(\frac{\partial}{\partial J} \right) \ln J e^{Jt} \Bigg|_{J=0} \\ &= f(t) + n f \left(\frac{\partial}{\partial J} \right) \int_0^{\infty} \frac{ds}{s} (e^{-s} - e^{-sJ}) e^{Jt} \Bigg|_{J=0} \\ &= f(t) + n \int_0^{\infty} \frac{ds}{s} [e^{-s} f(t) - f(t-s)]. \quad (\text{A5}) \end{aligned}$$

By the change of variable $s = e^l$ Eq. (A4) follows.

Appendix B: Asymptotic behaviour of the function B_r

In this appendix we study the asymptotic behavior for large λ of the function $B_r(\lambda x)$. By definition in Eq. (47)

$$\frac{1}{\lambda^r} B_r(\lambda x) := \sum_{k=1}^{\infty} (-1)^{k-1} \frac{e^{\lambda x k}}{(\lambda k)^r (k!)^2}, \quad (\text{B1})$$

so that

$$\begin{aligned} \frac{B_r(\lambda x)}{\lambda^r} &= -\frac{1}{\Gamma(r)} \int_0^{\infty} t^{r-1} \sum_{k=1}^{\infty} \frac{(-1)^k}{(k!)^2} e^{\lambda(x-t)k} dt \\ &= -\frac{1}{\Gamma(r)} \int_0^{\infty} t^{r-1} \left\{ J_0 \left[2e^{\frac{1}{2}\lambda(x-t)} \right] - 1 \right\} dt \\ &\xrightarrow{\lambda \rightarrow +\infty} \frac{1}{\Gamma(r)} \int_0^{\infty} t^{r-1} \theta(x-t) dt = \frac{x^r}{\Gamma(r+1)} \theta(x), \quad (\text{B2}) \end{aligned}$$

where we have used the fact that

$$J_0(x) := \sum_{m=0}^{\infty} \frac{(-1)^m}{(m!)^2} \left(\frac{x}{2} \right)^{2m} = \begin{cases} 1 & \text{when } x \rightarrow 0, \\ 0 & \text{when } x \rightarrow +\infty, \end{cases} \quad (\text{B3})$$

is the Bessel function of zero-th order of the first kind.

Appendix C: Some properties of the function \hat{G}_r

In the present Appendix we give some properties of the function \hat{G}_r , defined by the integral equation (12c). From the definition, we have that, for $0 \leq \alpha < \beta + 1$ and $r > -1$,

$$\begin{aligned} \hat{G}_r^{(\alpha)}(l) &:= \mathcal{D}_l^{\alpha} \hat{G}_r(l) = \\ &= \int_{-\infty}^{+\infty} \frac{(l+y)^{r-\alpha}}{\Gamma(r-\alpha+1)} e^{-\hat{G}_r(y)} \theta(l+y) dy. \quad (\text{C1}) \end{aligned}$$

Observe that

$$\hat{G}_r^{(\alpha)}(l) \geq 0 \quad \text{for } 0 \leq \alpha < r + 1. \quad (\text{C2})$$

In the previous equation we have used the fact that, for $0 \leq \alpha < \beta + 1$, we have [32]

$$\mathcal{D}_t^\alpha \left[\frac{t^\beta}{\Gamma(\beta + 1)} \theta(t) \right] = \frac{t^{\beta-\alpha}}{\Gamma(\beta - \alpha + 1)} \theta(t). \quad (\text{C3})$$

In particular, for $\alpha = r$ we have the simple relation

$$\hat{G}_r^{(r)}(l) := \mathcal{D}_l^r \hat{G}_r(l) = \int_{-\infty}^{\infty} e^{-\hat{G}_r(y)} \theta(y+l) dy. \quad (\text{C4})$$

Moreover, for $0 \leq \alpha < r + 1$,

$$\lim_{l \rightarrow -\infty} \hat{G}_r^{(\alpha)}(l) = 0. \quad (\text{C5})$$

From Eq. (C4)

$$\hat{G}_r^{(r+1)}(l) = e^{-\hat{G}_r(-l)} \geq 0 \Rightarrow \lim_{l \rightarrow +\infty} \hat{G}_r^{(r+1)}(l) = 1. \quad (\text{C6})$$

The relations above imply that

$$J_r^{(\alpha)} := \int_{-\infty}^{+\infty} \hat{G}_r(-u) \mathcal{D}_u^\alpha \hat{G}_r(u) du > 0, \quad 0 \leq \alpha < r + 1. \quad (\text{C7})$$

Similarly, for $0 < k < r + 1$ integer,

$$\begin{aligned} J_r^{(r+k+1)} &:= \int_{-\infty}^{+\infty} \hat{G}_r(-u) \mathcal{D}_u^{r+k+1} \hat{G}_r(u) du \\ &= \int_{-\infty}^{+\infty} \hat{G}_r(-u) \frac{d^k}{du^k} e^{-\hat{G}_r(-u)} du \\ &= \int_{-\infty}^{+\infty} \hat{G}_r^{(k)}(u) e^{-\hat{G}_r(u)} du \geq 0. \end{aligned} \quad (\text{C8})$$

For large l we have

$$\hat{G}_r(l) \approx \frac{l^{r+1}}{\Gamma(r+2)} \quad (\text{C9})$$

$$\hat{G}_r(-l) \approx \exp \left[-\frac{l^{r+1}}{\Gamma(r+2)} \right]. \quad (\text{C10})$$

As anticipated, an exact solution is available in the $r = 0$ case. In particular, for $r = 0$, the second derivative

$$\hat{G}_0^{(2)}(l) = e^{-\hat{G}_0(-l)} \hat{G}_0^{(1)}(-l) = \hat{G}_0^{(1)}(l) \hat{G}_0^{(1)}(-l) \quad (\text{C11})$$

is an even function of l

$$\hat{G}_0^{(2)}(l) - \hat{G}_0^{(2)}(-l) = 0 \Rightarrow \hat{G}_0^{(1)}(l) + \hat{G}_0^{(1)}(-l) = c \quad (\text{C12})$$

with the constant $c = 1$ by evaluating the left hand side in the limit of infinite l , and

$$\hat{G}_0^{(1)}(0) = e^{-\hat{G}_0(0)} = \frac{1}{2}. \quad (\text{C13})$$

Then we have that

$$\hat{G}_0(l) - \hat{G}_0(-l) = l \Rightarrow \hat{G}_0^{(1)}(l) = e^{-\hat{G}_0(-l)} = e^{l - \hat{G}_0(l)} \quad (\text{C14})$$

which means that

$$\frac{d}{dl} e^{\hat{G}_0(l)} = e^l \Rightarrow e^{\hat{G}_0(x)} - e^{\hat{G}_0(0)} = e^x - 1 \quad (\text{C15})$$

where we have used the the initial condition at $l = 0$, that is, because of Eq. (C13),

$$e^{\hat{G}_0(x)} = 1 + e^x \Rightarrow \hat{G}_0(x) = \ln(1 + e^x). \quad (\text{C16})$$

Appendix D: Evaluation of the integrals in the $r = 0$ case

To explicitly evaluate some of the integrals above, let us introduce the *polygamma function*

$$\psi_m(z) := \frac{d^{m+1}}{dz^{m+1}} \ln \Gamma(z) = (-1)^{m+1} \int_0^{\infty} \frac{t^m e^{-zt}}{1 - e^{-t}} dt, \quad (\text{D1})$$

which satisfies the recursion relation

$$\psi_m(z+1) = \psi_m(z) + (-1)^m \frac{m!}{z^{m+1}} \quad (\text{D2})$$

that, for a positive integer argument and assuming $m \geq 1$, leads to

$$\frac{\psi_m(k)}{(-1)^{m+1} m!} = \zeta(m+1) - \sum_{r=1}^{k-1} \frac{1}{r^{m+1}} = \sum_{r=k}^{\infty} \frac{1}{r^{m+1}}. \quad (\text{D3})$$

For $m = 0$ this implies

$$\psi_0(k) = -\gamma_E + H_{k-1} \Rightarrow \psi_0(1) = -\gamma_E, \quad (\text{D4})$$

with γ_E is Euler's gamma constant and

$$H_n := \sum_{k=1}^n \frac{1}{k} \quad (\text{D5})$$

are the harmonic numbers.

With this considerations in mind, and using Eq. (C16), we have that

$$\begin{aligned} J_0^{(1)} &:= \int_{-\infty}^{+\infty} \frac{\ln(1 + e^y)}{1 + e^y} dy \\ &= \int_0^{+\infty} \frac{t e^{-t}}{1 - e^{-t}} dt = \psi_1(1) = \zeta(2) = \sum_{k \geq 1} \frac{1}{k^2} = \frac{\pi^2}{6}. \end{aligned} \quad (\text{D6})$$

Afterwards we compute

$$\begin{aligned}
J_0^{(0)} &:= \int_{-\infty}^{+\infty} dy \frac{1}{1+e^y} \int_{-\infty}^y du \ln(1+e^u) \\
&= \int_0^{+\infty} dt \frac{e^{-t}}{1-e^{-t}} \int_0^t dw \frac{w}{1-e^{-w}} = - \int_0^{+\infty} dt t \frac{\ln(1-e^{-t})}{1-e^{-t}} \\
&= \sum_{k=1}^{\infty} \frac{1}{k} \int_0^{+\infty} dt \frac{te^{-kt}}{1-e^{-t}} = \sum_{k=1}^{\infty} \frac{1}{k} \psi_1(k). \quad (\text{D7})
\end{aligned}$$

We remark now that

$$\begin{aligned}
\sum_{k \geq 1} \frac{\psi_1(k)}{k} &= \sum_{k=1}^{\infty} \sum_{r=0}^{\infty} \frac{1}{k} \frac{1}{(r+k)^2} \\
&= \sum_{s \geq 1} \sum_{k=1}^s \frac{1}{k} \frac{1}{s^2} = \sum_{s \geq 1} \frac{1}{s^2} H_s. \quad (\text{D8})
\end{aligned}$$

Applying now the identity

$$\sum_{s=1}^{\infty} \frac{H_s}{s^2} = 2\zeta(3), \quad (\text{D9})$$

discovered by Euler, we recover the result obtained by Parisi and Ratiéville [25]

$$J_0^{(0)} = 2\zeta(3) = -\psi_2(1). \quad (\text{D10})$$

To finally evaluate $J_0^{(3)}$, we remark now that

$$\int_{-\infty}^{+\infty} \frac{dy}{1+e^y} \frac{d}{dy} \ln(1+e^y) = - \int_{-\infty}^{+\infty} dy \frac{d}{dy} \frac{1}{1+e^y} = 1. \quad (\text{D11})$$

Then, as

$$\frac{d^2}{dy^2} \ln(1+e^y) = \frac{d}{dy} \ln(1+e^y) - \left[\frac{d}{dy} \ln(1+e^y) \right]^2, \quad (\text{D12})$$

we have

$$\begin{aligned}
J_0^{(3)} &= \int_{-\infty}^{+\infty} dy \frac{1}{1+e^y} \frac{d^2}{dy^2} \ln(1+e^y) = \\
&= - \int_{-\infty}^{+\infty} dy \left(\frac{d}{dy} \frac{1}{1+e^y} \right) \frac{d}{dy} \ln(1+e^y) \\
&= \int_{-\infty}^{+\infty} dy \frac{1}{1+e^y} \left[\frac{d}{dy} \ln(1+e^y) \right]^2 = \frac{1}{2}. \quad (\text{D13})
\end{aligned}$$

Appendix E: Calculation of ΔF_r^T

To evaluate explicitly ΔF_r^T , let us start from Eq. (63),

$$\begin{aligned}
\Delta \hat{F}_r^T &= \frac{1}{2\hat{\beta}N} \sum_{s=1}^{\infty} \sum_{t=1}^{\infty} (-1)^{s+t-1} \frac{(s+t-1)!}{s!t!} \frac{\hat{g}_s \hat{g}_t}{\hat{g}_{s+t}^2} q_{s+t}^2 = \frac{\eta_0 \Gamma(r+1)}{2\hat{\beta}^{r+2}N} \sum_{s=1}^{\infty} \sum_{t=1}^{\infty} \frac{(-1)^{s+t-1}}{s!t!} \left(\frac{t+s}{st} \right)^{r+1} \int_{-\infty}^{+\infty} e^{-G_r(y)} e^{y(s+t)} q_{s+t} dy \\
&= \frac{\eta_0 \Gamma(r+1)}{2\hat{\beta}^{r+2}N} \sum_{k=2}^{\infty} \sum_{s=1}^{k-1} \frac{(-1)^{k-1}}{s!(k-s)!} \frac{k^{r+1}}{s^{r+1}(k-s)^{r+1}} \int_{-\infty}^{+\infty} e^{-G_r(y)} e^{yk} q_k dy, \quad (\text{E1})
\end{aligned}$$

and, in order to perform the sum over s , we introduce integral representations

$$\sum_{s=1}^{k-1} \frac{1}{s!(k-s)!} \frac{1}{s^{r+1}(k-s)^{r+1}} = \sum_{s=1}^{k-1} \binom{k}{s} \int_0^{+\infty} du \int_0^{+\infty} dv \frac{u^r v^r e^{-su} e^{-(k-s)v}}{k! \Gamma^2(r+1)} = \int_0^{+\infty} du \int_0^{+\infty} dv u^r v^r \frac{(e^{-u} + e^{-v})^k - e^{-uk} - e^{-vk}}{k! \Gamma^2(r+1)}. \quad (\text{E2})$$

Observing now that the value $k=1$ can be included in the sum over k , and denoting by

$$h := \frac{\hat{\beta}}{[\eta_0 \Gamma(r+1)]^{1/(r+1)}} \quad (\text{E3})$$

we can write

$$\begin{aligned}
\frac{2\hat{\beta}^{r+2}\Gamma(r+1)N}{\eta_0}\Delta\hat{F}_r^T &= \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k!} q_k \int_{-\infty}^{+\infty} dy e^{-G_r(y)} \mathcal{D}_y^{r+1} e^{yk} \int_0^{+\infty} du \int_0^{+\infty} dv u^r v^r [(e^{-u} + e^{-v})^k - e^{-uk} - e^{-vk}] \\
&= \int_{-\infty}^{+\infty} dy e^{-G_r(y)} \mathcal{D}_y^{r+1} \left\{ \int_0^{+\infty} du \int_0^{+\infty} dv u^r v^r [G_r(y + \ln(e^{-u} + e^{-v})) - G_r(y - u) - G_r(y - v)] \right\} \\
&= 2h^r \int_{-\infty}^{+\infty} dy e^{-\hat{G}_r(y)} \mathcal{D}_y^{r+1} \left\{ \int_0^{+\infty} du \int_0^u dv u^r v^r [G_r(h(y - v) + \ln(e^{-h(u-v)} + 1)) - \hat{G}_r(y - u) - \hat{G}_r(y - v)] \right\}. \quad (E4)
\end{aligned}$$

This implies that, for $h \rightarrow \infty$,

$$\begin{aligned}
\Delta\hat{F}_r^T &= \frac{1}{N\eta_0^{\frac{1}{r+1}}\Gamma^{2+\frac{1}{r+1}}(r+1)} \int_{-\infty}^{+\infty} dy e^{-\hat{G}_r(y)} \mathcal{D}_y^{r+1} \left[\int_0^{+\infty} du \int_0^u dv u^r v^r \hat{G}_r(y - u) \right] \\
&= -\frac{1}{N\eta_0^{\frac{1}{r+1}}\Gamma^{2+\frac{1}{r+1}}(r+1)} \frac{1}{r+1} \int_{-\infty}^{+\infty} dy e^{-\hat{G}_r(y)} \mathcal{D}_y^{r+1} \left[\int_{-\infty}^y (y-u)^{2r+1} \hat{G}_r(u) du \right] \\
&= -\frac{\Gamma(2r+2)}{N\eta_0^{\frac{1}{r+1}}\Gamma^{3+\frac{1}{r+1}}(r+1)} \frac{1}{r+1} \int_{-\infty}^{+\infty} dy e^{-\hat{G}_r(y)} \int_{-\infty}^y (y-u)^r \hat{G}_r(u) du = -\frac{\Gamma(2r+2)}{N\eta_0^{\frac{1}{r+1}}\Gamma^{2+\frac{1}{r+1}}(r+1)} \frac{1}{r+1} \int_{-\infty}^{+\infty} \hat{G}_r(-u)\hat{G}_r(u) du, \quad (E5)
\end{aligned}$$

that is exactly Eq. (64).

-
- [1] C. H. Papadimitriou and K. Steiglitz, *Combinatorial optimization: algorithms and complexity* (Courier Dover Publications, 1998).
- [2] L. Lovász and D. Plummer, *Matching Theory*, AMS Chelsea Publishing Series (2009).
- [3] H. W. Kuhn, *Naval Res. Logistics Quart.* **2**, 83 (1955).
- [4] J. Edmonds, *Can. J. Math.* **17**, 449 (1965).
- [5] J. Edmonds and R. M. Karp, *J. ACM* **19**, 248 (1972).
- [6] S. Micali and V. V. Vazirani, in *21st Annual Symposium on Foundations of Computer Science* (1980) pp. 17–27.
- [7] J. Steele, *Probability Theory and Combinatorial Optimization*, CBMS-NSF Regional Conference Series in Applied Mathematics (SIAM, 1997).
- [8] M. Mézard and G. Parisi, *J. Phys. Lett.* **46**, 771 (1985).
- [9] P. W. Shor, *Random planar matching and bin packing*, PhD thesis, MIT (1985).
- [10] M. Mézard and G. Parisi, *J. Phys.* **49**, 2019 (1988).
- [11] J. Yukich, *Probability theory of classical Euclidean optimization problems*, Lecture notes in mathematics (Springer, 1998).
- [12] C. Lucibello, G. Parisi, and G. Sicuro, *Phys. Rev. E* **95**, 012302 (2017).
- [13] S. Caracciolo, C. Lucibello, G. Parisi, and G. Sicuro, *Phys. Rev. E* **90**, 012118 (2014).
- [14] E. Boniolo, S. Caracciolo, and A. Sportiello, *J. Stat. Mech.* **2014**, P11023 (2014).
- [15] S. Caracciolo and G. Sicuro, *Phys. Rev. E* **91**, 062125 (2015).
- [16] S. Caracciolo and G. Sicuro, *Phys. Rev. E* **90**, 042112 (2014).
- [17] S. Caracciolo and G. Sicuro, *Phys. Rev. Lett.* **115**, 230601 (2015).
- [18] L. Ambrosio, F. Stra, and D. Trevisan, *ArXiv* (2016), [arXiv:1611.04960](https://arxiv.org/abs/1611.04960).
- [19] G. Sicuro, *The Euclidean matching problem*, Springer theses (Springer, 2017).
- [20] S. Kirkpatrick, C. Gelatt, and M. Vecchi, *Science* **220**, 671 (1983).
- [21] H. Orland, *J. Phys. Lett.* **46**, 763 (1985).
- [22] M. Mézard and A. Montanari, *Information, Physics, and Computation*, Oxford Graduate Texts (OUP Oxford, 2009).
- [23] V. Bapst, L. Foini, F. Krzakala, G. Semerjian, and F. Zamponi, *Phys. Rep.* **523**, 127 (2013).
- [24] M. Mézard and G. Parisi, *J. Phys.* **48**, 1451 (1987).
- [25] G. Parisi and M. Ratiéville, *Eur. Phys. J. B* **29**, 457 (2002).
- [26] S. F. Edwards and P. W. Anderson, *Journal of Physics F: Metal Physics* **5**, 965 (1975).
- [27] D. J. Aldous, *Random Struct. Algorithms*, 381 (2001).
- [28] G. Parisi, *ArXiv* (1998), [arXiv:9801176](https://arxiv.org/abs/9801176) [cond-mat].
- [29] S. Linusson and J. Wästlund, *Probab. Theory Relat. Fields* **128**, 419 (2004).
- [30] C. Nair, B. Prabhakar, and M. Sharma, *Random Struct. Algorithms* **27**, 413 (2005).
- [31] S. Samko, A. Kilbas, and O. Marichev, *Fractional Integrals and Derivatives* (Taylor & Francis, 1993).
- [32] I. Podlubny, *Fractional Differential Equations*, Mathematics in Science and Engineering (Academic Press, 1999).
- [33] J. Houdayer, J. H. Boutet de Monvel, and O. C. Martin, *Eur. Phys. J. B* **6**, 383 (1998).
- [34] G. Parisi and M. Ratiéville, *Eur. Phys. J. B* **22**, 229 (2001).
- [35] J. Salez, *Electron. Commun. Probab.* **20**, 7 pp. (2015).
- [36] R. Brunetti, W. Krauth, M. Mézard, and G. Parisi, *Europhys. Lett.* **14**, 295 (1991).

- [37] Y. Lee and J. B. Orlin, in *Large Scale Optimization: State of the Art*, edited by W. W. Hager, D. W. Hearn, and P. M. Pardalos (Kluwer Academic Publishers B.V., 1993) pp. 206–244.
- [38] R. Jonker and A. Volgenant, *Computing* **38**, 325 (1987).